



ISSN:2229-712X

Available online at [www.elixirjournal.org](http://www.elixirjournal.org)

## Applied Mathematics

Elixir Appl. Math. 42 (2012) 6369-6372

Elixir Online  
Journal

# Application of new transform "tarig transform" to partial differential equations

Tarig. M. Elzaki<sup>1</sup> and Salih M. Elzaki<sup>2</sup><sup>1</sup>Department of Mathematics, Faculty of Sciences and Arts-Alkamil, King Abdulaziz University, Jeddah Saudi Arabia<sup>2</sup>Department of Mathematics, Sudan University of Science and Technology.

### ARTICLE INFO

#### Article history:

Received: 22 October 2011;

Received in revised form:

15 January 2012;

Accepted: 25 January 2012;

### ABSTRACT

In this paper we derive the formulate for Tarig transform of partial derivatives and apply them to solve five types of initial value problems. Our purpose here is to show that the applicability of this interesting new transform and its effecting to solve such problems.

© 2012 Elixir All rights reserved.

### Keywords

Tarig Transform-

Partial Differential Equations.

### Introduction

The differential equations have played a central role in every aspect of applied mathematics for every long time and with the advent of the computer, their importance has increased father.

Thus investigation and analysis of differential equations cruising in applications led to many deep mathematical problems; therefore, there are so many different techniques in order to solve differential equations.

In order to solve the differential equations, the integral transforms were extensively used and thus there are several words on the theory and applications of integral transforms such as the Laplace, Fourier, Mellin, Hankel and Sumudu, to name but a few. Recently, Tarig M. Elzaki introduced a new integral transform, named Tarig transform, and further applied it to find the solution of ordinary and partial differential equations.

#### Definition and Derivations Tarig Transform of Derivatives

Tarig transform is defined as:

$$T[f(t)] = \frac{1}{u} \int_0^{\infty} f(t) e^{-\frac{t}{u^2}} dt = F(u), \quad t > 0, u \neq 0 \quad (1)$$

To obtain Tarig transform of Partial derivatives we use integration by parts as follows:

$$T\left[\frac{\partial f}{\partial t}(x,t)\right] = \frac{1}{u} \int_0^{\infty} \frac{\partial f}{\partial t} e^{-\frac{t}{u^2}} dt$$

Integrating by parts to find:

$$\begin{aligned} T\left[\frac{\partial f}{\partial t}\right] &= \frac{1}{u} \left[ \left[ f(x,t) e^{-\frac{t}{u^2}} \right]_0^{\infty} + \frac{1}{u^2} \int_0^{\infty} e^{-\frac{t}{u^2}} f(x,t) dt \right] \\ &= \frac{1}{u^2} T[f(x,t)] - \frac{1}{u} f(x,0) \end{aligned} \quad (2)$$

We assume that  $f$  is piece wise continuous and is of exponential order.

Now

$$\begin{aligned} T\left[\frac{\partial f}{\partial x}\right] &= \int_0^{\infty} \frac{1}{u} e^{-\frac{t}{u^2}} \frac{\partial f}{\partial x}(x,t) dt = \frac{\partial}{\partial x} \int_0^{\infty} \frac{1}{u} e^{-\frac{t}{u^2}} f(x,t) dt \\ &= \frac{d}{dx} [T(f(x,t))] = \frac{d}{dx} [F(u)] \end{aligned} \quad (3)$$

Also we can find that:

$$T\left[\frac{\partial^2 f}{\partial x^2}\right] = \frac{d^2}{dx^2} [F(u)] \quad (4)$$

To find:  $T\left[\frac{\partial^2 f}{\partial t^2}(x,t)\right]$  Let  $\frac{\partial f}{\partial t} = g$ , then,

$$T\left[\frac{\partial^2 f}{\partial t^2}(x,t)\right] = T\left[\frac{\partial g}{\partial t}(x,t)\right] = \frac{T[g(x,t)]}{u^2} - \frac{1}{u} g(x,0)$$

By equation (2) we have,

$$T\left[\frac{\partial^2 f}{\partial t^2}(x,t)\right] = \frac{1}{u^4} T[f(x,t)] - \frac{1}{u^3} f(x,0) - \frac{1}{u^2} \frac{\partial f}{\partial t}(x,0) \quad (5)$$

We can easily extend this result to the nth partial derivative by using mathematical induction.

#### Solution of Partial Differential Equations

In this paper we solve the linear first and second order partial differential equations, which are fundamental equations in mathematical physics' and occur in many branches of physics, a applied mathematics as well as in engineering.

#### Example 1:

Find the solution of the first order initial value problem,

$$\frac{\partial y}{\partial x} = 2 \frac{\partial y}{\partial t} + y, \quad y(x,0) = 6e^{-3x} \quad (6)$$

And  $y$  is bounded for  $x > 0$ ,  $t > 0$ .

Tele:

E-mail addresses: [Tarig.alzaki@gmail.com](mailto:Tarig.alzaki@gmail.com)

© 2012 Elixir All rights reserved

**Solution:**

Let  $F(u)$  be Tarig transform of, then, taking Tarig transform of (6) to get:

$$\frac{d}{dx} [F(u)] = 2 \left[ \frac{1}{u^2} F(u) - \frac{1}{u} y(x, 0) \right] + F(u)$$

Where  $F(u)$  is Tarig transform of  $y(t, x)$  using the initial condition to find that:

$$\frac{d}{dx} F(u) - \left( \frac{2}{u^2} + 1 \right) F(u) = -\frac{12}{u} e^{-3x}$$

This is the linear ordinary differential equation, the integrating

factor is  $e^{-\frac{(2+u^2)}{u^2}x}$  therefore:

$$F(u) = \frac{12}{u} \left[ \frac{u^2}{2+4u^2} \right] e^{-3x} + ce^{\frac{2+u^2}{u^2}x}$$

$F(u)$  is bounded, then  $c = 0$  and

$$F(u) = 6 \left[ \frac{u}{1+2u^2} \right] e^{-3x}$$

Taking the inverse Tarig transform to find:

$$y(x, t) = 6e^{-3x} e^{-2t} = 6e^{-3x-2t}$$

**Example 2:**

Consider the one dimensional unsteady heat conduction problem as follows:

$$\frac{\partial U}{\partial t} = \frac{\partial^2 U}{\partial x^2}, \quad t > 0, \quad 0 < x < 1 \quad (7)$$

With the boundary conditions:

$$U(x, 0) = 3 \sin 2\pi x, \quad U(0, t) = U(1, t) = 0 \quad (8)$$

**Solution:**

Taking Tarig transform of eq (7) we have:

$$\frac{F(x, u)}{u^2} - \frac{1}{u} U(x, 0) = \frac{d^2}{dx^2} F(x, u)$$

Or

$$u^2 D^2 F(x, u) - F(x, u) = -3u \sin 2\pi x \quad (9)$$

The solutions of (9) are,

$$F_c(x, u) = c_1 e^{\frac{x}{u}} + c_2 e^{-\frac{x}{u}},$$

$$F_p(x, u) = \frac{-3u \sin 2\pi x}{u^2 D^2 - 1} = \frac{3u \sin 2\pi x}{1 + 4\pi^2 u^2}$$

Then:

$$F(x, u) = c_1 e^{\frac{x}{u}} + c_2 e^{-\frac{x}{u}} + \frac{3u \sin 2\pi x}{1 + (2\pi)^2 u^2}$$

Taking Tarig transform of the boundary conditions,

$$T[U(0, t)] = F(0, u) = 0, \quad T[U(1, t)] = F(1, u) = 0,$$

Then we have:  $c_1 + c_2 = 0$  and  $ce^{\frac{1}{u}} + c_1 e^{-\frac{1}{u}} = 0$

These means that  $c_1 = c_2 = 0$ , then the solution of (9) is,

$$F(x, u) = \frac{3u \sin 2\pi x}{1 + 4\pi^2 u^2}$$

Then:

$$U(x, t) = F^{-1} \left[ \frac{3u \sin 2\pi x}{1 + 4\pi^2 u^2} \right] = 3e^{-4\pi^2 t} \sin 2\pi x \quad (*)$$

This problem has an interesting physical interpretation. If we consider a solid bounded by the infinite plane faces

$x = 0$  and  $x = 1$ , the equation.  $\frac{\partial U}{\partial t} = k \frac{\partial^2 U}{\partial x^2}$  is the

equation for heat conduction in this solid where  $U = U(x, t)$

is the temperature at any plane face  $x$  at any time  $t$  and  $k$  is a constant called the diffusivity, which depends on the material of the solid. The boundary conditions

$U(0, t) = 0$  and  $U(1, t) = 0$  Indicate that the

temperatures at  $x = 0$  and  $x = 1$  are kept at temperature

zero, while  $U(x, 0) = 3 \sin 2\pi x$  represents the initial

temperature everywhere in  $0 < x < 1$ , result (\*) is the

temperature Everywhere in the solid at time  $t > 0$ .

**Example 3:**

Consider the following wave equation,

$$\frac{\partial^2 w(x, t)}{\partial t^2} - 4 \frac{\partial^2 w(x, t)}{\partial x^2} = 0, \quad 0 \leq x \leq 1, \quad t > 0 \quad (10)$$

With the initial conditions:

$$w(x, 0) = \sin \pi x, \quad \frac{\partial w}{\partial t}(x, 0) = 0 \quad (11)$$

And the boundary conditions,

$$w(0, t) = w(1, t) = 0 \quad (12)$$

**Solution:**

Applying Tarig transform to eq(10) we have:

$$\frac{\bar{w}(x, u)}{u^4} - \frac{w(x, 0)}{u^3} - \frac{w_t(x, 0)}{u} - 4\bar{w}_{xx}(x, u) = 0 \quad (13)$$

Where  $\bar{w}(x, u)$  is Tarig transform of  $w(x, t)$ . substituting

eq(11) into eq(13) we get:

$$\bar{w}(x, u) - 4u^4 \bar{w}_{xx}(x, u) = u \sin \pi x \quad (14)$$

Like example (2), using the boundary conditions (12) we get

that:  $w_c(x, u) = 0$

Then the solution of (14) is,

$$\bar{w}(x, u) = \frac{u \sin \pi x}{1 - 4u^4 D^2} = \frac{u \sin \pi x}{1 + (2\pi)^2 u^4},$$

Where  $D^2 = \frac{d^2}{dx^2}$

Then:

$$w(x,t) = \sin \pi x F^{-1} \left[ \frac{u}{1+(2\pi)^2 u^4} \right] = \sin \pi x \cos 2\pi x$$

**Example 4:**

Consider the following Laplace equation,

$$U_{xx}(x,y) + U_{yy}(x,y) = 0, \quad t > 0 \quad (15)$$

With the initial conditions:

$$U(x,0) = 0, \quad U_y(x,0) = \cos x \quad (16)$$

And the boundary conditions:

$$U(0,t) = U(1,t) = 0, \quad 0 \leq x \leq 1 \quad (17)$$

**Solution:**

By using Tarig transform into eq(15) yields:

$$\bar{U}_{xx}(x,u) + \frac{\bar{U}(x,u)}{u^4} - \frac{U(x,0)}{u^3} - \frac{U_y(x,0)}{u} = 0 \quad (18)$$

Using the initial conditions (16) to find:

$$u^4 \bar{U}_{xx}(x,u) + \bar{U}(x,u) = u^3 \cos x \quad (19)$$

The solutions of equation (19) under the boundary conditions (17) are,

$$\bar{U}_c(x,u) = 0 \quad \text{and} \quad \bar{U}_p = \frac{u^3 \cos x}{1-u^4}$$

Then the general solution of (15) is,

$$U(x,y) = \cos x F^{-1} \left[ \frac{u^3}{1-u^4} \right] = \cos x \sinh y$$

**Example 5:**

Consider the following Telegrapher's equation,

$$U_{tt}(x,t) + 2\alpha U_t(x,t) = \alpha^2 U_{xx}(x,t), \quad 0 < x < 1, \quad t > 0 \quad (20)$$

Where  $\alpha$  is positive constant.

With the initial conditions:

$$U(x,0) = \cos x, \quad U_t(x,0) = 0 \quad (21)$$

And the boundary conditions:

$$U(0,t) = 0, \quad U(1,t) = 0 \quad (22)$$

**Solution:**

Using Tarig transform into eq(20) and making use of the initial conditions (21) we have:

$$\alpha^2 u^4 \bar{U}_{xx}(x,u) - (1+2\alpha u^2) \bar{U}(x,u) = -u \cos x - 2\alpha u^3 \cos x$$

Taking Tarig transform of the boundary conditions (22) we get:

$$\bar{U}(0,u) = 0, \quad \bar{U}(1,u) = 0$$

Use these conditions to obtain the general solution of eq (23) in the form:

$$\bar{U}(x,u) = \frac{u + 2\alpha u^3 \cos x}{1 + 2\alpha u^2 - \alpha^2 u^4 D^2}, \quad \left( D^2 = \frac{d^2}{dx^2} \right)$$

$$\bar{U}(x,u) = u \cos x \left[ \frac{1+2\alpha u^2}{(1+\alpha u^2)^2} \right] = \cos x \left[ \frac{u}{1+\alpha u^2} + \frac{\alpha u^3}{(1+\alpha u^2)^2} \right]$$

Then:

$$U(x,t) = \cos x F^{-1} \left[ \frac{u}{1+\alpha u^2} + \frac{\alpha u^3}{(1+\alpha u^2)^2} \right] = \cos x [e^{-\alpha t} + \alpha t e^{-\alpha t}]$$

And,  $U(x,t) = (1 + \alpha t) e^{-\alpha t} \cos x$

**Conclusion**

Application of tarig transform to solution of special different partial differential equation has been demonstrated.

**References:**

- [1] Tarig M. Elzaki, The New Integral Transform "Elzaki Transform" Global Journal of Pure and Applied Mathematics, ISSN 0973-1768, Number 1(2011), pp. 57-64.
- [2] Tarig M. Elzaki & Salih M. Elzaki, Application of New Transform "Elzaki Transform" to Partial Differential Equations, Global Journal of Pure and Applied Mathematics, ISSN 0973-1768, Number 1(2011), pp. 65-70.
- [3] Tarig M. Elzaki & Salih M. Elzaki, On the Connections Between Laplace and Elzaki transforms, Advances in Theoretical and Applied Mathematics, ISSN 0973-4554 Volume 6, Number 1(2011), pp. 1-11.
- [4] Tarig M. Elzaki & Salih M. Elzaki, On the Elzaki Transform and Ordinary Differential Equation With Variable Coefficients, Advances in Theoretical and Applied Mathematics. ISSN 0973-4554 Volume 6, Number 1(2011), pp. 13-18.
- [5] Tarig M. Elzaki, Adem Kilicman, Hassan Eltayeb. On Existence and Uniqueness of Generalized Solutions for a Mixed-Type Differential Equation, Journal of Mathematics Research, Vol. 2, No. 4 (2010) pp. 88-92.
- [6] Tarig M. Elzaki, Existence and Uniqueness of Solutions for Composite Type Equation, Journal of Science and Technology, (2009). pp. 214-219.
- [7] Tarig M. Elzaki, and Salih M. Elzaki, On the new integral Transform 'Tarig Transform' and Systems of Ordinary Differential Equations. Applied Mathematics, Elixir Appl. Math. 36 (2011) 3226-3229.
- [8] Tarig M. Elzaki, and Salih M. Elzaki, On the relationship between Laplace Transform and new integral transform Tarig Transform'. Applied Mathematics, Elixir Appl. Math. 36 (2011) 3230-3233.
- [9] Tarig M. Elzaki, and Salih M. Elzaki, On the Tarig Transform and ordinary differential equation with variable coefficients, Applied Mathematics, Elixir Appl. Math. 38 (2011) 4250-4252.
- [10] Tarig M. Elzaki, and Salih M. Elzaki, the new integral transform "Tarig Transform" Properties and applications to differential equations, Applied Mathematics, Elixir Appl. Math. 38 (2011) 4239-4242.
- [11] Lokenath Debnath and D. Bhatta. Integral transform and their Application second Edition, Chapman & Hall /CRC (2006).
- [12] A.Kilicman and H.E.Gadain. An application of double Laplace transform and Sumudu transform, Lobachevskii J. Math.30 (3) (2009), pp.214-223.
- [13] J. Zhang, Asumudu based algorithm m for solving differential equations, Comp. Sci. J. Moldova 15(3) (2007), pp - 303-313.

[14] Hassan Eltayeb and Adem kilicman, A Note on the Sumudu Transforms and differential Equations, Applied Mathematical Sciences, VOL, 4,2010, no.22,1089-1098.

[15] Kilicman A. & H. ELtayeb. A note on Integral transform and Partial Differential Equation, Applied Mathematical Sciences, 4(3) (2010), PP.109-118.

[16] Hassan ELtayeh and Adem kilicman, on Some Applications of a new Integral Transform, Int. Journal of Math. Analysis, Vol, 4, 2010, no.3, 123-132.

**Appendix**  
**Tarig Transform of Some Functions**

S.NO.	$f(t)$	$F(u)$
1	1	$u$
2	$t$	$u^3$
3	$e^{at}$	$\frac{u}{1-au^2}$
4	$t^n$	$n! u^{2n+1}$
5	$t^a$	$\Gamma(a+1)u^{2a+1}$
6	$\sin at$	$\frac{au^3}{1+a^2u^4}$
7	$\cos at$	$\frac{u}{1+a^2u^4}$
8	$\sinh at$	$\frac{au^3}{1-a^2u^4}$
9	$\cosh at$	$\frac{u}{1-a^2u^4}$
10	$H(t-a)$	$ue^{\frac{-a}{u^2}}$
11	$\delta(t-a)$	$\frac{1}{u}e^{\frac{-a}{u^2}}$
12	$te^{at}$	$\frac{u^3}{(1-au^2)^2}$